

DEFORMATIONS OF ASSOCIATIVE ALGEBRAS WITH INNER PRODUCTS

JOHN TERILLA AND THOMAS TRADLER

ABSTRACT. We develop the deformation theory of A_∞ algebras together with ∞ -inner products and identify a differential graded Lie algebra that controls the theory. This generalizes the deformation theories of associative algebras, A_∞ algebras, associative algebras with inner products, and A_∞ algebras with inner products.

CONTENTS

1. Introduction	1
2. Definitions of A_∞ algebras and ∞ inner products	2
2.1. A_∞ algebras	2
2.2. A_∞ bimodules.	3
2.3. ∞ inner products	3
2.4. Induced maps	3
3. Deformations of A_∞ algebras and ∞ inner products	4
3.1. Deformations of A_∞ algebras	5
3.2. Deformations of A_∞ algebras with ∞ inner products	6
4. Moduli, infinitesimal deformations, and relationship to cyclic cohomology	8
Appendix A. Explicit formulas of $\delta_f(i)$	9
References	14

1. INTRODUCTION

In this paper, we describe a differential graded Lie algebra that controls deformations of the homotopy versions of associative algebras together with the inner products introduced in [9]. Let us review the basic idea of a deformation theory governed by a differential graded Lie algebra [7, 2, 3, 4].

Fix a ground field k of characteristic $\neq 2$. For any differential graded Lie (dg Lie) algebra $(\mathfrak{g} = \oplus_i \mathfrak{g}^i, d, [,]) over k , one can consider deforming the differential d in the direction of an inner derivation. Informally, such a deformation is given by an (equivalence classes of) α making$

$$d_\alpha := d + \text{ad}(\alpha)$$

into a differential. The map d_α is always a derivation and the condition that $d_\alpha^2 = 0$ translates into the Maurer-Cartan equation:

$$d\alpha + \frac{1}{2}[\alpha, \alpha] = 0.$$

Key words and phrases. homotopy, inner product, deformation theory.

The authors would like to thank Jim Stasheff, Martin Markl, and Dennis Sullivan for many helpful discussions.

The deformed differential d_α may involve parameters from the maximal ideal m of a $\mathbb{Z}$ graded Artin local ring: $\alpha \in (\mathfrak{g} \otimes_k m)^1$. If m is the maximal ideal of a local Artin ring R and $\alpha \in (\mathfrak{g} \otimes_k m)^1$ is a solution to the Maurer-Cartan equation, then one may call d_α a *deformation of d over R* . Two deformations of d should be considered equivalent if they are related via a natural automorphism of $\mathfrak{g} \otimes m$. A ring map $R \rightarrow S$ will transport a deformation of d over R to a deformation of d over S .

More formally, one has a functor $Def_{\mathfrak{g}}$ from the category of $\mathbb{Z}$ graded Artin local rings with residue field k to the category of sets, assigning to such a ring R with maximal ideal m the set

$$Def_{\mathfrak{g}}(R) = \{\alpha \in (\mathfrak{g} \otimes_k m)^1 : d\alpha + \frac{1}{2}[\alpha, \alpha] = 0\} / \sim.$$

Here, $\sim$ is the equivalence relation determined by the action of the gauge group, which we now recall. Since R is an Artin ring, m is a nilpotent algebra, and $(\mathfrak{g} \otimes_k m)^0 \subseteq \mathfrak{g} \otimes_k m$ is a nilpotent Lie algebra. Therefore, there exists a group $G = \{\exp \beta : \beta \in (\mathfrak{g} \otimes_k m)^0\}$, called the gauge group, with multiplication defined by the Baker-Campbell-Hausdorff formula. The action of $e^\beta \in G$ on an element $\alpha \in (\mathfrak{g} \otimes_k m)^1$ is determined by the infinitesimal action:

$$\alpha \mapsto \beta \cdot \alpha = [\beta, \alpha] - d\beta, \quad \alpha \in (\mathfrak{g} \otimes m)^1, \beta \in (\mathfrak{g} \otimes m)^0.$$

This action satisfies

$$e^{\text{ad } \beta} d_\alpha e^{-\text{ad } \beta} = d_{e^{\beta} \cdot \alpha},$$

and preserves the set of solutions to the Maurer-Cartan equation. In this paper, we work with A_∞ algebras equipped with an ∞ inner product. One has the notion of a deformation of an A_∞ algebra with an ∞ inner product over a ring R , and there is a natural equivalence on the set of deformations. A ring map $R \rightarrow S$ transports deformations over R to deformations over S . The association

$$R \mapsto \left\{ \begin{array}{c} \text{deformations of the } A_\infty \text{ algebra with} \\ \text{the } \infty \text{ inner product over } R \end{array} \right\} / \left\{ \begin{array}{c} \text{equivalent} \\ \text{deformations} \end{array} \right\}$$

defines a covariant deformation functor.

In this paper, we construct a $\text{dgLa}(\mathfrak{h} = \oplus_i \mathfrak{h}^i, d, [\cdot, \cdot])$ associated to an A_∞ algebra with an ∞ inner product, and prove that the functor described above is isomorphic to $Def_{\mathfrak{h}}$. This is the precise mathematical content of the statement *the dgLa $(\mathfrak{h}, [\cdot, \cdot], d)$ controls the deformations of the A_∞ algebra with an ∞ inner product*.

2. DEFINITIONS OF A_∞ ALGEBRAS AND ∞ INNER PRODUCTS

We now review the concept of an ∞ inner product on an A_∞ algebra [9], [8]. The concepts of A_∞ algebras, A_∞ bimodules, A_∞ bimodule maps, and A_∞ inner products are generalizations of the usual concepts of associative algebras, bimodules, bimodule maps, and invariant inner products.

2.1. A_∞ algebras. Let $V = \bigoplus_{j \in \mathbb{Z}} V^j$ be a graded module over a ring S . Recall that the suspension $V[1]$ of V is defined to be $V[1] = \bigoplus_{j \in \mathbb{Z}} (V[1])^j$ with $(V[1])^j := V^{j-1}$. For a graded S -module A , BA denotes the tensor coalgebra $BA := T(A[1])$, sometimes called the *bar complex* of A . An A_∞ algebra over S is defined to be a pair (A, D) where A is a graded S module and $D \in \text{Coder}(BA)$ of degree -1 with $D^2 = 0$.

Suppose that (A, D) and (A', D') are A_∞ algebras over S . Then an A_∞ map from (A', D') to (A, D) is a map $\lambda : BA' \rightarrow BA$ satisfying $\lambda \circ D' = D \circ \lambda$.

2.2. A_∞ bimodules. Let (A, D) be an A_∞ algebra over S , and let M be a graded S module. Let $B^M A$ denote the tensor bicomodule $B^M A := T^{M[1]}(A[1]) = \bigoplus_{k,l \geq 0} A[1]^{\otimes k} \otimes M[1] \otimes A[1]^{\otimes l}$ of $M[1]$ over BA . An A_∞ *bimodule structure on M over A* is defined to be a coderivation $D^M \in \text{Coder}_D(B^M A, B^M A)$ over D of degree -1 with $(D^M)^2 = 0$.

Let (M, D^M) and (N, D^N) be A_∞ bimodules over A . Let $\text{Comap}(B^M A, B^N A)$ denote the maps $F : B^M A \rightarrow B^N A$ satisfying

$$\begin{array}{ccc} B^M A & \xrightarrow{\Delta^M} & (BA \otimes B^M A) \oplus (B^M A \otimes BA) \\ \downarrow F & & \downarrow (\text{Id} \otimes F) \oplus (F \otimes \text{Id}) \\ B^N A & \xrightarrow{\Delta^N} & (BA \otimes B^M A) \oplus (B^M A \otimes BA) \end{array}$$

The space $\text{Comap}(B^M A, B^N A)$ carries a differential defined by

$$\delta^{M,N}(F) := D^N \circ F - (-1)^{|F|} F \circ D^M.$$

In this case, an A_∞ *bimodule map from M to N* is defined to be an element $F \in \text{Comap}(B^M A, B^N A)$ of degree 0 with $\delta^{M,N}(F) = 0$, i.e.

$$D^N \circ F = F \circ D^M.$$

2.3. ∞ inner products. For any $f \in \text{Coder}(BA)$, there are induced coderivations $f^A \in \text{Coder}_f(B^A A, B^A A)$ and $f^{A^*} \in \text{Coder}_f(B^{A^*} A, B^{A^*} A)$, where $A^* = \text{hom}_S(A, S)$ denotes the dual of A . One also has an induced map

$$\delta_f : \text{Comap}(B^M A, B^N A) \rightarrow \text{Comap}(B^M A, B^N A)$$

given by $\delta_f(F) = f^{A^*} \circ F - (-1)^{|f||F|} \cdot F \circ f^A$. Note that, in particular, if (A, D) is an A_∞ algebra, then A and A^* have A_∞ bimodule structures given by D^A and D^{A^*} .

Definition 2.1. Let (A, D) be an A_∞ algebra over S . We define an ∞ *inner product on A over S* to be an A_∞ bimodule map I from A to A^* . Equivalently, an ∞ inner product is an element $I \in \text{Comap}(B^A A, B^{A^*} A)$ satisfying

$$\delta_D(I) = D^{A^*} \circ I - I \circ D^A = 0.$$

Every inner product $\langle, \rangle : A \otimes A \rightarrow S$ defines an element $I \in \text{Comap}(B^A A, B^{A^*} A)$. In this case, the condition $D^{A^*} \circ I - I \circ D^A = 0$ is equivalent to $\langle D(a_1, \dots, a_n), a_{n+1} \rangle = \pm \langle a_1, D(a_2, \dots, a_{n+1}) \rangle$. See the appendix for additional illustrations.

2.4. Induced maps. Recall, that if $\lambda : A' \rightarrow A$ is an algebra map between two associative algebras, then every module over A is also a module over A' , and similarly for module maps. Also, $\lambda : A' \rightarrow A$ and $\lambda^* : A^* \rightarrow (A')^*$ will be module maps over A' . Here we give the corresponding homotopy generalizations.

Suppose that λ is an A_∞ map from (A', D') to (A, D) . First, every A_∞ bimodule (M, D^M) over A is also an A_∞ bimodule over A' , whose structure map is determined by the lowest components (which are maps $B^M A \rightarrow M$)

$$\begin{aligned} (D^M)^\lambda(a'_1, \dots, a'_k, m, a'_{k+1}, \dots, a'_{k+l}) \\ = \sum \pm pr_M \circ D^M(\lambda(a'_1, \dots), \dots, \lambda(\dots, a'_k), m, \lambda(a'_{k+1}, \dots), \dots, \lambda(\dots, a'_{k+l})). \end{aligned}$$

Here, pr_M denotes the projection onto M . The signs are given by the usual sign rule, namely introducing a sign $(-1)^{|\alpha| \cdot |\beta|}$, whenever α jumps over β . The relevant degrees are the degrees given in $B^M A$.

Also, any A_∞ bimodule map $F : B^M A \rightarrow B^N A$ over A induces an A_∞ bimodule map $F^\lambda : B^M A' \rightarrow B^N A'$ over A' given by

$$\begin{aligned} F^\lambda(a'_1, \dots, a'_k, m, a'_{k+1}, \dots, a'_{k+l}) \\ = \sum \pm pr_N \circ F(\lambda(a'_1, \dots), \dots, \lambda(\dots, a'_k), m, \lambda(a'_{k+1}, \dots), \dots, \lambda(\dots, a'_{k+l})). \end{aligned}$$

Furthermore λ induces the two A_∞ bimodule maps over A'

$$\bar{\lambda} : B^{A'} A' \rightarrow B^A A' \text{ and } \tilde{\lambda} : B^{A^*} A' \rightarrow B^{(A')^*} A'$$

defined by the components

$$\bar{\lambda}(a'_1, \dots, a'_{k+l+1}) = pr_A \circ \lambda(a'_1, \dots, a'_{k+l+1})$$

and

$$(\tilde{\lambda}(a'_1, \dots, a'^*, \dots, a'_{k+l}))(a') = \pm a'^*(pr_A \circ \lambda(a'_{k+1}, \dots, a'_{k+l}, a', a'_1, \dots, a'_k)).$$

For our own comfort, we write TA and $T^M A$ instead of BA and $B^M A$ for the remainder of this paper.

3. DEFORMATIONS OF A_∞ ALGEBRAS AND ∞ INNER PRODUCTS

Before we define the specific differential graded Lie algebra $(\mathfrak{h}, d, [, \cdot])$ that controls the deformations of A_∞ structures and ∞ inner products, we discuss a simple example and make a remark.

Example 3.1. Any graded associative algebra $\mathfrak{g}$ becomes a Lie algebra by defining the bracket to be the usual commutator. An element $\alpha \in \mathfrak{g}^1$ satisfying $\alpha^2 = 0$ is sometimes called a *polarization*. With a polarization $\alpha \in \mathfrak{g}^1$, $\mathfrak{g}$ becomes a differential graded Lie algebra by setting the differential to be $\delta = \text{ad}(\alpha)$. With δ so defined, the Mauer-Cartan equation becomes

$$0 = \delta(\gamma) + \frac{1}{2}[\gamma, \gamma] = \frac{1}{2}[\alpha + \gamma, \alpha + \gamma].$$

In other words, $\gamma \in \mathfrak{g}^1$ satisfies the Mauer-Cartan equation if and only if $\alpha + \gamma$ is another polarization.

Now let S be a graded ring and consider $\mathfrak{g}$ defined by

$$\mathfrak{g} = \left\{ \begin{pmatrix} a & 0 \\ b & a \end{pmatrix} \middle| a, b \in S \right\}$$

with the bracket defined as the usual graded commutator of matrix multiplication:

$$\left[\begin{pmatrix} a & 0 \\ b & a \end{pmatrix}, \begin{pmatrix} c & 0 \\ d & c \end{pmatrix} \right] = \begin{pmatrix} [a, c] & 0 \\ [b, c] + [a, d] & [a, c] \end{pmatrix}.$$

Then, $\begin{pmatrix} D & 0 \\ I & D \end{pmatrix} \in \mathfrak{g}^1$ is a polarization if and only if

$$0 = [D, D] = 2 \cdot D^2 \text{ and } 0 = [D, I] + [I, D] = 2 \cdot [D, I].$$

Having chosen a polarization $P = \begin{pmatrix} D & 0 \\ I & D \end{pmatrix}$, the formula for $\delta = \text{ad}(P)$ is given by

$$\delta \begin{pmatrix} f & 0 \\ i & f \end{pmatrix} = \left[\begin{pmatrix} D & 0 \\ I & D \end{pmatrix}, \begin{pmatrix} f & 0 \\ i & f \end{pmatrix} \right] = \begin{pmatrix} [D, f] & 0 \\ [D, i] + [f, I] & [D, f] \end{pmatrix}.$$

Now we look at the gauge equivalence. First of all, the gauge group $G = \exp(\mathfrak{g}^0)$ is the Lie group consisting of matrices of the form e^A , for any $A \in \mathfrak{g}^0$. The gauge action of G on $\mathfrak{g}$ is then determined by $e^{\text{ad}(A)} \cdot B = \text{Ad}(e^A)(B) = e^A B e^{-A}$. A computation shows that

$$\exp \begin{pmatrix} f & 0 \\ i & f \end{pmatrix} = \begin{pmatrix} e^f & 0 \\ x & e^f \end{pmatrix}, \text{ where } x = \sum_{n \geq 1} \frac{1}{n!} \sum_{k+l=n-1} f^k \cdot i \cdot f^l.$$

Then the gauge equivalence summarizes as

$$e^A \begin{pmatrix} D & 0 \\ I & D \end{pmatrix} e^{-A} = \begin{pmatrix} e^f D e^{-f} & 0 \\ e^f I e^{-f} + [e^f D e^{-f}, x e^{-f}] & e^f D e^{-f} \end{pmatrix}.$$

This concludes the example.

Remark 3.2. Let N be a graded coalgebra over S . Then $\text{hom}(N, N)$ will be a graded associative algebra and a Lie algebra with the bracket defined by the commutator. The space $\text{Coder}(N) \subseteq \text{hom}(N, N)$ will not, in general, form an associative algebra, but it is a Lie algebra. In particular, for any vector space A , $\text{Coder}(TA)$ is a graded Lie algebra. An A_∞ structure on A consists of an element $D \in \text{Coder}(TA)$ satisfying $D^2 = 0$. Thus one can say that an A_∞ structure on A is a choice of polarization $D \in \text{Coder}(TA)$. Hence, if (A, D) is an A_∞ algebra, $\text{Coder}(TA)$ carries a differential $\delta : \text{Coder}(TA) \rightarrow \text{Coder}(TA)$ defined by

$$\delta(f) := [D, f] = D \circ f - (-1)^{|f|} f \circ D.$$

The complex $(\text{Coder}(TA), \delta)$ is called the Hochschild cochain complex of A . Together with the bracket from $\text{hom}(TA, TA)$, it is a differential graded Lie algebra that controls the deformations of the A_∞ algebra (A, D) . In order to make this statement precise, we recall the deformation theory of A_∞ algebras (see for example [1]). As a first observation, one may note that γ is a solution to the Maurer-Cartan equation in the Hochschild dgLa if and only if $D + \gamma$ is another polarization in $\text{Coder}(TA)$; i.e., another A_∞ structure, on A .

3.1. Deformations of A_∞ algebras. Let A be a graded vector space over a field k and let R be a graded Artin local algebra with residue field k . Let m denote the maximal ideal of R . We have the decomposition $R \simeq R/m \oplus m \simeq k \oplus m$ and the projection $pr_k : R \rightarrow k$, hence the decomposition $A \otimes R \simeq A \oplus (A \otimes m)$ and the projection $pr_A : A \otimes R \rightarrow A$. For definiteness, the reader may have the concrete example $R = k[t]/t^{l+1}$ in mind. In this example, the maximal ideal is $m = tk[t]/t^{l+1}$, $A \otimes R \simeq A + At + At^2 + \cdots + At^l$ (with the tensor signs suppressed) and the natural projection pr_A maps $a_0 + a_1 t + a_2 t^2 + \cdots + a_l t^l \mapsto a_0$.

Let (A, D) be an A_∞ algebra over k . A *deformation of (A, D) over R* is an A_∞ algebra $(A \otimes R, D')$ over R with the property that the projection

$$pr : T(A \otimes R) \simeq TA \otimes R \rightarrow TA$$

is a morphism of A_∞ algebras over k . This means that $pr \circ D' = D \circ pr$.

Suppose that D' is a deformation of (A, D) over R . Via any map $R \rightarrow S$, one can view $A \otimes R$ as an S module and $(A \otimes R, D')$ as a deformation of (A, D) over S .

Let $\pi \in \text{hom}(R \otimes R, R)$ denote the multiplication in R . Let D_R denote the A_∞ structure $D \otimes \pi$ on $A \otimes R$. The A_∞ algebra $(A \otimes R, D_R)$ is the model for a trivial deformation of (A, D) . That is, $(A \otimes R, D')$ is a *trivial deformation* if it is isomorphic to $(A \otimes R, D_R)$ as an A_∞ algebra. This means that there is an automorphism

$$\lambda : T(A \otimes R) \rightarrow T(A \otimes R)$$

satisfying $\lambda \circ D' = D_R \circ \lambda$. Two deformations are equivalent if and only if they differ by a trivial one.

3.2. Deformations of A_∞ algebras with ∞ inner products.

Definition 3.3. Let A be a graded vector space over a field k . We define the graded Lie algebra $(\mathfrak{h} = \oplus_i \mathfrak{h}^i, [\cdot, \cdot])$ by

$$(1) \quad \mathfrak{h}^i = \text{Coder}(TA)^{-i} \oplus \text{Comap}(T^A A, T^{A^*} A)^{1-i}$$

and

$$(2) \quad \begin{aligned} [(f, i), (g, j)] &= ([f, g], \delta_f(j) - (-1)^{|f||g|} \delta_g(i)) \\ &= (fg - (-1)^{|f||g|} gf, f^{A^*} j - (-1)^{|f||j|} j f^A - (-1)^{|f||g|} g^{A^*} i + (-1)^{|g| \cdot (|f| + |i|)} i g^A). \end{aligned}$$

The skew-symmetry and Jacobi identity of $[\cdot, \cdot]$ are straightforward to check after one notices that $\delta_f \circ \delta_g - (-1)^{|f||g|} \delta_g \circ \delta_f = \delta_{f \circ g} - (-1)^{|f||g|} g \circ f$.

Proposition 3.4. A pair $(D, I) \in \mathfrak{h}$ is an A_∞ structure with ∞ inner product on A if and only if $[(D, I), (D, I)] = 0$.

Proof. This is immediate:

$$0 = [(D, I), (D, I)] \Leftrightarrow 0 = [D, D] = 2 \cdot D^2 \text{ and } 0 = 2 \cdot \delta_D(I) = 2(D^{A^*} \circ I - I \circ D^A).$$

The condition $D^2 = 0$ means that D defines an A_∞ structure on A and the condition $D^{A^*} \circ I - I \circ D^A = 0$ means that I defines a compatible ∞ -inner product. $\square$

Now fix an A_∞ structure together with an ∞ inner product, which is to say, fix a pair $(D, I) \in \mathfrak{h}$ with $[(D, I), (D, I)] = 0$. Then, define $d : \mathfrak{h} \rightarrow \mathfrak{h}$ by

$$(3) \quad d(f, i) = [(D, I), (f, i)].$$

The triple $(\mathfrak{h}, d, [\cdot, \cdot])$ is a differential graded Lie algebra.

Definition 3.5. A deformation of an A_∞ algebra with ∞ inner product (A, D, I) over R is an A_∞ algebra over R with ∞ inner product $(A \otimes R, D', I')$, such that the projection

$$pr : T(A \otimes R) \rightarrow TA$$

is a morphism of A_∞ algebras over k compatible with the ∞ -inner products. Compatibility with the ∞ inner product means that the following diagram of A_∞ -bimodule maps over k is commutative:

$$\begin{array}{ccc} T^{A \otimes R}(A \otimes R) & \xrightarrow{\overline{pr}} & T^A(A \otimes R) \\ \downarrow pr_k \circ I' & & \downarrow I^{pr} \\ T^{(A \otimes R)^*}(A \otimes R) & \xleftarrow{\overline{pr}} & T^{A^*}(A \otimes R) \end{array}$$

Here, the ∞ -inner product I' on $A \otimes R$ over R induces an ∞ -inner product on $A \otimes R$ over k by composing with the map induced by the projection $\text{hom}_R(A \otimes R, R) \rightarrow \text{hom}_k(A \otimes R, k)$, $f \mapsto pr_k \circ f$.

There is a natural extension of I to an ∞ -inner product $I_R = I \otimes \pi$ on $(A \otimes R, D_R)$.

Definition 3.6. We say that (D', I') is a *trivial deformation* of (D, I) provided “the triple $(A \otimes R, D', I')$ is isomorphic to $(A \otimes R, D_R, I_R)$ as A_∞ algebras with ∞ inner products.” That is, if there exist an automorphism

$$\lambda : T(A \otimes R) \rightarrow T(A \otimes R)$$

and a comap

$$\rho : T^{A \otimes R}(A \otimes R) \rightarrow T^{(A \otimes R)^*}(A \otimes R)$$

satisfying

- (i) $\lambda \circ D' = D_R \circ \lambda$,
- (ii) $I' - \tilde{\lambda} \circ (I_R)^\lambda \circ \bar{\lambda} = D'^{(A \otimes R)^*} \circ \rho + \rho \circ D'^{A \otimes R}$.

It may be helpful to think of the second condition in definition 3.6 as saying I' equals I_R under a change of coordinates (given by λ) up to a homotopy (given by ρ). That is, the following diagram commutes, up to a homotopy defined by $\rho \in \text{Comap}(T^{A \otimes R}(A \otimes R))$.

$$\begin{array}{ccc}
 T^{A \otimes R}(A \otimes R) & \xrightarrow{\bar{\lambda}} & T^{A \otimes R}(A \otimes R) \\
 \downarrow I' & & \downarrow (I_R)^\lambda \\
 T^{(A \otimes R)^*}(A \otimes R) & \xleftarrow{\tilde{\lambda}} & T^{(A \otimes R)^*}(A \otimes R)
 \end{array}$$

Two deformations are equivalent if and only if they differ by a trivial one.

Now, the conclusion:

Theorem 3.7. *Let (A, D) be an A_∞ algebra and let I be an ∞ -inner product. Then the dg Lie algebra $(\mathfrak{h}, d, [\cdot, \cdot])$ defined by equations (1), (2) and (3) controls the deformations of the A_∞ algebra with ∞ inner product (A, D, I) .*

Proof. The content of this theorem is summarized in the following two statements.

- Deformations, over R , of the (A, D, I) correspond to solutions to the Maurer-Cartan equation in $\mathfrak{h} \otimes m$,
- and equivalent deformations correspond to gauge equivalent solutions.

First we prove the first statement. Let $\alpha = (f, i) \in (\mathfrak{h} \otimes m)^1$. Observe that

$$\begin{aligned}
 d\alpha + \frac{1}{2}[\alpha, \alpha] &= [(D_R, I_R), (f, i)] + \frac{1}{2} \cdot [(f, i), (f, i)] \\
 &= \frac{1}{2} \cdot [(D_R + f, I_R + i), (D_R + f, I_R + i)].
 \end{aligned}$$

Then, proposition 3.4 proves that $d\alpha + \frac{1}{2}[\alpha, \alpha] = 0$ if and only if $(A \otimes R, D_R + f, I_R + i)$ is a deformation of (A, D, I) . It is immediate that any $(A \otimes R, D', I')$ that is a deformation of (A, D, I) must satisfy $[(D', I'), (D', I')] = 0 \in \mathfrak{h} \otimes R$. The fact that $pr : T(A \otimes R) \rightarrow TA$ is a map of A_∞ algebras with ∞ inner products implies that $D' = D_R + f$ and $I' = I_R + i$ for some $(f, i) \in \mathfrak{h} \otimes m$.

Now we prove the second statement. Let $\alpha = (f, i) \in (\mathfrak{h} \otimes m)^0$. The gauge action for $\mathfrak{h}$ becomes

$$e^{\text{ad}(f, i)} \cdot (D_R, I_R) = \sum_{n \geq 0} \frac{\text{ad}(f, i)^n}{n!} (D_R, I_R).$$

It follows from

$$\delta_f (\delta_{\text{ad}(f)^r(D_R)}((\delta_f)^s(i))) = \delta_{\text{ad}(f)^{r+1}(D_R)}((\delta_f)^s(i)) + \delta_{\text{ad}(f)^r(D_R)}((\delta_f)^{s+1}(i)),$$

that $\text{ad}(f, i)^n(D_R, I_R)$ is given by

$$\left(\text{ad}(f)^n(D_R), (\delta_f)^n(I_R) - \sum_{k+l=n-1} \frac{n!}{k!(l+1)!} \cdot \delta_{\text{ad}(f)^k(D_R)} \circ (\delta_f)^l(i) \right).$$

Now define $\lambda^{-1} = e^f = \sum_{k \geq 0} \frac{1}{k!} f^k$ and $\rho = \sum_{l \geq 0} \frac{-1}{(l+1)!} \cdot (\delta_f)^l(i)$. Then for the automorphism λ and the homotopy ρ , we have

$$\begin{aligned} & \sum_{n \geq 0} \frac{\text{ad}(f, i)^n}{n!} (D_R, I_R) \\ &= \left(\sum_{n \geq 0} \frac{\text{ad}(f)^n}{n!} (D_R), \sum_{n \geq 0} \frac{(\delta_f)^n}{n!} (I_R) + \delta_{\sum_{k \geq 0} \frac{\text{ad}(f)^k}{k!} (D_R)} \left(\sum_{l \geq 0} \frac{-1}{(l+1)!} \cdot (\delta_f)^l(i) \right) \right) \\ &= \left(\lambda^{-1} D_R \lambda, \tilde{\lambda}(I_R)^\lambda \bar{\lambda} + \delta_{\lambda^{-1} D_R \lambda}(\rho) \right). \end{aligned}$$

This proves that $e^{\text{ad}(f, i)} \cdot (D_R, I_R)$ is a trivial deformation of (D, I) .

It is not hard to see that every trivial deformation of (D, I) arises from an element gauge equivalent to the identity. The condition that the A_∞ algebra map $\lambda : T(A \otimes R) \rightarrow T(A \otimes R)$ is an automorphism implies that $\lambda = e^f$ for some $f \in (\text{Coder}(TA) \otimes m)^0$. Also, since $\rho = -i - \frac{1}{2} \delta_f(i) \cdots$, the map $i \mapsto \rho(i) = \sum_{l \geq 0} \frac{-1}{(l+1)!} \cdot (\delta_f)^l(i)$ is invertible. So one can obtain any homotopy ρ , by choosing a suitable element $i = \sum_{m \geq 0} c_m \cdot (\delta_f)^m(\rho) \in (\mathfrak{h} \otimes m)^0$ with $\rho(i) = \rho$. $\square$

4. MODULI, INFINITESIMAL DEFORMATIONS, AND RELATIONSHIP TO CYCLIC COHOMOLOGY

Let us return briefly to general deformation theory in order to review the notions of infinitesimal deformations and moduli space. Let $(\mathfrak{g}, d, [, \cdot])$ be a differential graded Lie algebra and assume that $\text{Ker}(d)/\text{Im}(d) =: H(\mathfrak{g}) = \bigoplus_{i=-m}^m H^i(\mathfrak{g})$ is finite dimensional. Consider the (graded version of the) ring of dual numbers $R = k[t_{-m}, \dots, t_m]/t_i t_j$. Here $\deg(t_i) = i - 1$ and the maximal ideal of R is $m = \bigoplus_i t_i R$.

From a solution $\sum(\gamma_j \otimes t_j) \in (\mathfrak{g} \otimes m)^1$ to the Maurer-Cartan equation, one may produce the map $d + \sum t_j \text{ad}(\gamma_j) : \mathfrak{g} \otimes k[t_{-m}, \dots, t_m] \rightarrow \mathfrak{g} \otimes k[t_{-m}, \dots, t_m]$ which satisfies

$$\left(d + \sum t_j \text{ad}(\gamma_j) \right)^2 = 0 \text{ modulo } t_i t_j.$$

One refers to $\gamma = \sum \gamma_j$ as an infinitesimal deformation. One can readily check that

$$\text{Def}_{\mathfrak{g}}(R) = \text{Ker}(d)/\text{Im}(d) = H(\mathfrak{g}).$$

Schlessinger's theorem [6] implies that $\text{Def}_{\mathfrak{g}}$ is prorepresentable. That is, there exists a projective limit of (graded) local Artin rings $\mathcal{O}$ and an equivalence of the functors

$$\text{Def}_{\mathfrak{g}}(\cdot) \simeq \text{hom}(\mathcal{O}, \cdot).$$

In the case that $\mathcal{O} = \mathcal{O}_{\mathcal{M}}$ is the ring of local functions at the base point of a pointed $\mathbb{Z}$ graded space $\mathcal{M}$, then $\mathcal{M}$ is the local moduli space for $\text{Def}_{\mathfrak{g}}$. Denote the base point of $\mathcal{M}$ by 0. One can check that

$$T_0(\mathcal{M}) \simeq \text{hom}(\mathcal{O}_{\mathcal{M}}, R).$$

It follows that the graded tangent space to the moduli space at the base point is isomorphic to the cohomology of $(\mathfrak{g}, d)$:

$$T_0(\mathcal{M}) \simeq H(\mathfrak{g}).$$

Now, let (A, D) be an A_∞ algebra and let I be an ∞ inner product on (A, D) . Theorem 3.7 says that the dg Lie algebra controlling deformations of (A, D, I) is

$$\mathfrak{h} = \text{Coder}(TA) \oplus \text{Comap}(T^A A, T^{A^*} A)$$

with bracket

$$[(f, i), (g, j)] = ([f, g], \delta_f(j) - (-1)^{|f||g|} \delta_g(i))$$

and a differential

$$d(f, i) = [(D, I), (f, i)].$$

Thus follows the expected infinitesimal statement:

Corollary 4.1. *The tangent space to the moduli space of A_∞ structures with ∞ inner products is isomorphic to $H(\mathfrak{h})$.*

As a final remark, we mention some connections between the cohomology $H(\mathfrak{h})$ and a couple of its cousins. If (A, D, I) is an A_∞ algebra with ∞ -inner product, we have the Hochschild dgLa $(\text{Coder}(TA), \delta, [,])$ and the sub dgLa of cyclic Hochschild cochains $\text{Coder}(TA)_{\text{Cyclic}}$, defined by

$$\text{Coder}(TA)_{\text{Cyclic}} = \{f \in \text{Coder}(TA) : \delta_f(I) = 0\}.$$

Note from appendix A that if I consists of an ordinary symmetric inner product $I = \langle \cdot, \cdot \rangle$, then the condition $\delta_f(I) = f^{A^*} \circ I - I \circ f^A = 0$ is equivalent to

$$\langle f(a_1, \dots, a_n), a_{n+1} \rangle = \pm \langle a_1, f(a_2, \dots, a_{n+1}) \rangle.$$

We have the following maps of differential graded Lie algebras.

$$(4) \quad (\text{Coder}(TA)_{\text{Cyclic}}, \delta, [,]) \longrightarrow (\mathfrak{h}, d, [,]) \text{ and } (\mathfrak{h}, d, [,]) \longrightarrow (\text{Coder}(TA), \delta, [,]).$$


The first map is the injection $f \mapsto (f, 0) \in \mathfrak{h}$, which is a cochain map


$$d(f, 0) = ([D, f], \pm(f^{A^*} \circ I - I \circ f^A)) = (\delta f, 0),$$

because elements of the domain are cyclic. The induced map in cohomology describes a statement from [5], namely that the first order deformations of D compatible with the inner product are classified by cyclic cohomology. We do not know under what conditions the map $f \mapsto (f, 0) \in \mathfrak{h}$ induces an isomorphism in cohomology. The second map in (4) is simply the projection $\text{Coder}(TA) \oplus \text{Comap}(T^A A, T^{A^*} A) \rightarrow \text{Coder}(TA)$ and the induced map in cohomology describes the simple statement that any infinitesimal deformation of the pair (D, I) gives an infinitesimal deformation of D .


APPENDIX A. EXPLICIT FORMULAS OF $\delta_f(i)$

Let $f \in \text{Coder}(TA)$ and $i \in \text{Comap}(T^A A, T^{A^*} A)$. We want to describe the term $\delta_f(i) = f^{A^*} \circ i - (-1)^{|f||i|} \cdot i \circ f^A \in \text{Comap}(T^A A, T^{A^*} A)$ more explicitly. Here, $f : \oplus_{k \geq 1} A^{\otimes k} \rightarrow A$ and $i : \oplus_{k, l \geq 0} A^{\otimes k} \otimes A \otimes A^{\otimes l} \otimes A \rightarrow S$ have the components


$$f_k : A^{\otimes k} \rightarrow A$$


$$i_{k,l} = \langle \cdots \rangle_{k,l} : A^{\otimes k} \otimes A \otimes A^{\otimes l} \otimes A \rightarrow S$$



By convention, the inputs are always inserted using the counterclockwise direction. Then $f^{A^*} \circ i - (-1)^{|f||i|} \cdot i \circ f^A$ is given by inserting f into i in all possible combinations.




First, here are some examples of how these diagrams are to be read.
 $\langle a, b, c, d \rangle_{2,0}$




$\langle a, b, c, d, e, f, g, h, i \rangle_{3,4}$




$$\langle f_2(f_2(b, c), f_2(d, e)), f_2(f, a) \rangle_{0,0}$$



$$\langle a, b, f_3(c, d, f_2(e, f)), g, f_2(h, i) \rangle_{1,2}$$



$$\langle c, f_2(d, e), f_2(f_2(f, g), h), i, f_4(j, k, a, b) \rangle_{2,1}$$



Here are the terms of $\delta_f(i) = f^{A^*} \circ i - (-1)^{|f||i|} \cdot i \circ f^A$ up to sign, when they are being applied to elements from $A^{\otimes k} \otimes A \otimes A^{\otimes l} \otimes A$:

$k = 0, l = 0$:

$$\langle f_1(a), b \rangle_{0,0} \pm \langle a, f_1(b) \rangle_{0,0}$$

$$a \text{---} \bullet \text{---} \circ \text{---} b \quad \pm \quad a \text{---} \circ \text{---} \bullet \text{---} b$$

$k = 1, l = 0$:

$$\begin{aligned} &\langle f_1(a), b, c \rangle_{1,0} \pm \langle a, f_1(b), c \rangle_{1,0} \pm \langle a, b, f_1(c) \rangle_{1,0} \pm \\ &\quad \langle f_2(a, b), c \rangle_{0,0} \pm \langle b, f_2(c, a) \rangle_{0,0} \end{aligned}$$

$$\begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \circ \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \circ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm$$

$$\begin{array}{c} a \\ \diagup \\ \bullet \\ | \\ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ \diagup \\ \bullet \\ | \\ \text{---} \circ \text{---} \bullet \text{---} c \\ | \\ b \end{array}$$

$k = 0, l = 1$:

$$\begin{aligned}
& \langle f_1(a), b, c \rangle_{0,1} \pm \langle a, f_1(b), c \rangle_{0,1} \pm \langle a, b, f_1(c) \rangle_{0,1} \pm \\
& \langle f_2(a, b), c \rangle_{0,0} \pm \langle a, f_2(b, c) \rangle_{0,0}
\end{aligned}$$

$$\begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \circ \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ | \\ \bullet \\ | \\ \text{---} \circ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm$$


$$\begin{array}{c} a \\ \diagup \\ \bullet \\ | \\ \text{---} \bullet \text{---} c \\ | \\ b \end{array} \pm \begin{array}{c} a \\ \diagup \\ \bullet \\ | \\ \text{---} \circ \text{---} \bullet \text{---} c \\ | \\ b \end{array}$$

$k = 2, l = 0$:

$$\begin{aligned}
& \langle f_1(a), b, c, d \rangle_{2,0} \pm \langle a, f_1(b), c, d \rangle_{2,0} \pm \\
& \langle a, b, f_1(c), d \rangle_{2,0} \pm \langle a, b, c, f_1(d) \rangle_{2,0} \pm \\
& \langle f_2(a, b), c, d \rangle_{1,0} \pm \langle a, f_2(b, c), d \rangle_{1,0} \pm \langle b, c, f_2(d, a) \rangle_{1,0} \pm \\
& \langle f_3(a, b, c), d \rangle_{0,0} \pm \langle c, f_3(d, a, b) \rangle_{0,0}
\end{aligned}$$

Note that for example the term $\langle a, b, f_2(c, d) \rangle_{2,0}$ does not appear, because c and d are the two special elements of $a \otimes b \otimes c \otimes d \in A^{\otimes 2} \otimes A \otimes A^{\otimes 0} \otimes A$, which are put on the horizontal line of the diagram. The two special elements from $A^{\otimes k} \otimes A \otimes A^{\otimes l} \otimes A$ can never be inside


any f_n .



$k = 0, l = 2$:

$$\begin{aligned} & \langle f_1(a), b, c, d \rangle_{0,2} \pm \langle a, f_1(b), c, d \rangle_{0,2} \pm \\ & \langle a, b, f_1(c), d \rangle_{0,2} \pm \langle a, b, c, f_1(d) \rangle_{0,2} \pm \\ & \langle f_2(a, b), c, d \rangle_{0,1} \pm \langle a, f_2(b, c), d \rangle_{0,1} \pm \langle a, b, f_2(c, d) \rangle_{0,1} \pm \\ & \langle f_3(a, b, c), d \rangle_{0,0} \pm \langle a, f_3(b, c, d) \rangle_{0,0} \end{aligned}$$


The special elements are a and d from $a \otimes b \otimes c \otimes d \in A^{\otimes 0} \otimes A \otimes A^{\otimes 2} \otimes A$.



$k = 1, l = 1$:

$$\begin{aligned} & \langle f_1(a), b, c, d \rangle_{1,1} \pm \langle a, f_1(b), c, d \rangle_{1,1} \pm \\ & \langle a, b, f_1(c), d \rangle_{1,1} \pm \langle a, b, c, f_1(d) \rangle_{1,1} \pm \\ & \langle f_2(a, b), c, d \rangle_{0,1} \pm \langle b, c, f_2(d, a) \rangle_{0,1} \pm \\ & \langle a, f_2(b, c), d \rangle_{1,0} \pm \langle a, b, f_2(c, d) \rangle_{1,0} \pm \\ & \langle f_3(a, b, c), d \rangle_{0,0} \pm \langle b, f_3(c, d, a) \rangle_{0,0} \end{aligned}$$

The special elements are b and d from $a \otimes b \otimes c \otimes d \in A^{\otimes 1} \otimes A \otimes A^{\otimes 1} \otimes A$.




$i = \langle , \rangle_{0,0}$ for any k, l : Assume that $i = \langle , \rangle_{0,0}$ has only lowest component, but f has all higher components. We apply $f^{A^*} \circ i - (-1)^{|f||i|} \cdot i \circ f^A$ to the element

$$a_1 \otimes \dots \otimes a_k \otimes a_{k+1} \otimes a_{k+2} \otimes \dots \otimes a_{k+l+1} \otimes a_{k+l+2} \in A^{\otimes k} \otimes A \otimes A^{\otimes l} \otimes A$$

to get

$$\langle f(a_1, \dots, a_{k+l+1}), a_{k+l+2} \rangle_{0,0} \pm \langle a_{k+1}, f(a_{k+2}, \dots, a_{k+l+2}, a_1, \dots, a_k) \rangle_{0,0}$$



REFERENCES

- [1] A. Fialowski and M. Penkava. Deformation theory of infinity algebras. [math.RT/0101097](#), 2001.
- [2] W.M. Goldman and J.J. Millson. The deformation theory of representations of fundamental groups of compact Kähler manifolds. In *Publ. Math IHES*, number 67, pages 43–96. IHES, 1988.
- [3] M. Kontsevich. Deformation quantization of Poisson manifolds, I. [math.QA/9709040](#), 1997. preprint.
- [4] M. Manetti. Deformation theory via differential graded Lie algebras. In *Seminari di Geometria Algebrica*, pages 21–48. Scuola Normale Superiore, 1999.
- [5] M. Penkava. Infinity algebras, cohomology and cyclic cohomology, and infinitesimal deformations. [math.QA/0111088](#), 2001.
- [6] M. Schlessinger. Functors of Artin rings. *Trans. Am. Math. Soc.*, (130):208–222, 1968.
- [7] M. Schlessinger and J. Stasheff. The Lie algebra structure of tangent cohomology and deformation theory. *J. Pure and Applied Algebra*, 38:313–322, 1985.
- [8] T. Tradler and M. Zeinalian. Poincaré duality at the chain level, 2002.
- [9] T. Tradler. Infinity inner products on A-infinity algebras. [math.AT/0108027](#), 2001.

E-mail address: jterilla@math.sunysb.edu

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, STONY BROOK, NY, 11794-3651.

E-mail address: tradler@stanford.edu

DEPARTMENT OF MATHEMATICS, BLDG. 380 STANFORD UNIVERSITY, STANFORD, CA 94305-2125.